



(index.html)

12' Frequency



What is AEB ?!

More details !

PrevNext

1

What is AEB ?!

More details !

- [Home \(index.php\)](#)
- [About](#)
- [About this journal](#)

Contact us for more info
([contact.html](#))

About AEB



AEB

Advances in Environmental Biology (AEB) is an international scientific journal edited and published by **AENSI** publications, Jordan. **(AEB)** exists to publish results of research in all area of **Environmental Biology**. Papers are welcomed from both developed and developing countries describing original investigations of interest to those engaged in scientific research. The material should not have been previously published elsewhere. Review articles on a particularly hot subject may be suggested to the editor-in-chief. Book reviews, news of international meetings and activities, letters to the editors and editorial notes will also be published at the editors' discretion The average time between submission and final decision is 30 days and the average time between acceptance and final publication is 32 days.

Contact Us



(index.html)

12' Frequency



Our Editorial Board..

a detailed view.

PrevNext

1

Our Editorial Board..

a detailed view.

- Home (index.php)
- Editorial Board

Contact us for more info
(contact.html)



Editor in Chief::

- Prof. Dr. Abdel Rahman Mohammad Said Al-Tawaha Department of Biological sciences, Al-Hussein bin Talal University, Maan, Jordan. Amman, Jordan, Tel.00962776693869 e-mail: abdeltawaha74@gmail.com

Co-Editor in Chief::

- Prof. Amanullah, Associate Professor of Agronomy, Department of Agronomy, Faculty of Crop Production Sciences, The University of Agriculture Peshawar, Pakistan, E-mail: amanullah@aup.edu.pk
- Prof. Hikmet Günal, Gaziosmanpaşa University, Faculty of Agriculture, Department of Soil Science and Plant Nutrition, Tokat, Turkey., E-mail: hikmetgunal@gmail.com
- Prof. Arun Karnwal, (Microbiology), School of Bioengineering and Biosciences, Lovely Professional University, Jalandhar-Delhi G.T. Road, National Highway 1, Phagwara,

Advisory Board::



- Dr. Ignacy Kitowski, Department of Nature Conservation, Institute of Biology, University of Maria-Curie Sklodowska, Akademicka 19, PL-20-033 Lublin, Poland.
- Dr. Andrzej Komosa, Department of Radiochemistry and Colloid Chemistry, Maria Curie Sklodowska University, Lublin, Poland.
- Dr. Robin DUPONNOIS, Directeur de Recherche à l'IRD, Laboratoire Commun de Microbiologie, IRD/ISRA/UCAD, Centre de Recherche de Bel-Air, BP 1386 CP 18524 Dakar-Sénégal
- Dr. Hidetaka HORI Ph.D., Laboratories of Applied Bioscience, Graduate School of Science and Technology, Niigata University, Niigata 950-2181, Japan
- Dr. Panos S. ECONOMIDIS, Professor Emeritus at the Aristotle University, Karakasi str. 79, GR-544 53 Thessaloniki, Greece
- Dr. Christopher (Kitt) E. Bagwell, Ph.D. , Savannah River National Laboratory, SRNL Environmental Sciences & Biotechnology
- Dr. Yuexia Wang, College of Bioscience and Biotechnology, Yangzhou University, NO. 12 East Wenhui Road, Yangzhou City, Jiangsu Province, China
- Dr. Marius Ciprian Branzila, Technical University of Iasi, Faculty of Electrical Engineering, Department of Electrical Measurements and Materials, Bd. Dimitrie Mangeron 53 Iasi, 700050 Romania

Editors::

- Dr. Cai zhiqian Xishuangbanna, Tropical Botanical Garden, The Chinese Academy of Sciences, Menglun, Yunnan 666303, P.R. China.
- Dr. F.M. Aminuzzaman, Department of Plant Pathology, Sher-e-Bangla Agricultural University, Sher-e-Bangla Nagar, Dhaka-1207, Bangladesh.
- Dr SSS Sarma, Professor & National Researcher, National Autonomous University of Mexico, Campus Iztacala AP 314, CP 54090 Los Reyes, Iztacala, Tlalnepantla Edo. de Mexico Mexico.
- Dr. YASIR HASAN SIDDIQUE FIBR, Human Genetics and Toxicology Lab, Section of Genetics, Department of Zoology, Faculty of Life Sciences, Aligarh Muslim University, Aligarh - 202002 (UP) India.
- Dr. P.R. Salve, Scientist, Environmental Impact and Risk Assessment Division, National Environmental Engineering Research Institute (NEERI), Nehru Marg, Nagpur-440 020(M.S.), India
- Dr. Nuray MISIR, Karadeniz Technical University, Faculty of Forestry, 61080, Trabzon, Turkey
- Dr. Bragadeeswaran CAS in Marine Biology, Annamalai University, Parangipettai 608 502 Cuddalore, Tamil Nadu, India
- Dr. Sevil TOROGLU Biology Department, Faculty of Arts and Sciences, University of KSU, 46100 Avsar Campus, Kahramanmaraş, Turkey
- Dr. Taiga Akpovughaye Department of Biological Sciences, K.S.U., P.M.B. 1008, Anyigba, Kogi State
- Dr. Ezekiel Olatunji Department of Fisheries, Cross River University of Technology, Calabar, P.M.B. 102, Obubra, Nigeria.
- Dr. Shahid A. Soomro Paul-Hindemith-Allee4/Apt. 312, D-80939 München, Germany
- Dr. YOUNGASPHREE NAIDOO Senior Lecturer, School of Biological and Conservation Sciences, UKZN –Westville campus
- Dr. Ranya Aly Helmy Amer Department of Environmental Biotechnology, Genetic Engineering and Biotechnology Research Institute (GEBRI), Mubarak City for Scientific Research and Technology Application, New Burg El-Arab City, Universities and Research Institutes Zone, 21934 Alexandria, Egypt
- Dr. Slavomír Čerňansk Comenius University in Bratislava, Faculty of Natural Sciences, Department of Ecosozology and Physiotactics, Mlynska dolina 1, 842 15 Bratislava, Slovakia
- Prof. Dr. Renato G. Reyes College of Arts and Sciences, Central Luzon State University, Science City of Muñoz, Nueva Ecija.
- Prof. Magdy Tawfik Khalil Zoology Dept., Fac. Science, Ain Shams Univ., Cairo, Egypt. Aquatic Ecology, Biodiversity, Management & Conservation
- Dr. Halil Erhan EROĞLU Department of Biology, 66200 YOZGAT, Bozok University, Faculty of Science and Arts, Turkey.
- Prof. Shikui DONG School of Environmental Sciences, Beijing Normal University, No. 19, Xingjiekou Waidajie, Aidian District, Beijing, 100875, P.R. China
- Dr. Naveed Ahmed Khan Senior Lecturer in Microbiology, School of Biological & Chemical Sciences, Birkbeck College, University of London, Malet Street, London WC1E 7HX, England, U.K.
- Dr. Ignacy Kitowski Department of Nature Conservation, Institute of Biology, Maria Curie-Sklodowska University, Akademicka 19, PL-20-033 Lublin, Poland
- Dr. Songhe Zhang College of Environment, Hohai University, XiKang road NO.1. Gulou district, Nanjing 210098, Jiangsu
- Dr. Oscar Martínez Alvarez Station de Biologie Marine du Muséum National d'Histoire, Naturelle et du Collège de France. BP 225, 29182 Concarneau Cedex, France.
- Dr. Piotr Tryjanowski Department of Behavioural Ecology, Adam Mickiewicz University, Umultowska 89, PL 61-614 Poznan, Poland
- Dr. Rafael Caballero García de Arévalo Centro de Ciencias Medioambientales, C/ Serrano 115 bis, Madrid 28006, Spain
- Dr. Ömür BAYSAL Turkish Ministry of Agriculture and Rural Affairs, West Mediterranean Agricultural Research Institute (BATEM), Plant Pathology Department, P.B. 35, 07100 Antalya/Turkey

- Dr. Li, Feng-Rui Department of Ecology and Agriculture, Cold and Arid Regions Environmental and Engineering Research Institute, Chinese Academy of Sciences, China
- Dr. Bob Redden Curator, Australian Temperate Field Crops Collection, DPI-Vic, Private mail bag 260, Horsham Vic 3401, Australia
- Dr. Ahmad K. Hegazy Head of Ecology Division, Prof. of Conservation & Applied Ecology, Botany Department, Faculty of Science, Cairo University, Giza 12613, Egypt
- Prof. Dr. Marcelo Enrique Conti SPES - Development Studies Research Centre, Università di Roma "La Sapienza" Via Del Castro Laurenziano 9, 00161 Rome, Italy
- Prof. Cesar G. Demayo, Professor of Genetics and Entomology, Department of Biological Sciences MSU-Iligan Institute of Technology ,9200 Iligan City, Philippines

Contact Us

-  Editor in Cheif - Dr. Abdel Rahman Tawaha
-  Address: Amman-Jordan
-  +962-795016606
-  Aensieditor@Gmail.Com (mailto:Aensieditor@Gmail.Com)
-  Jasaeditor@Gmail.Com (mailto:Jasaeditor@Gmail.Com)



[| Home](#) | [Journals](#) | [Special Issues](#) | [Conferences](#) | [Contact us](#) |

Special Issue IPN Kuching, 2015

Online Data Access Control with Bluetooth Beacon for Digital Content Stores

Young-Woong Ju and Min Choi

1-5

Investigation on Relationship between Human Behavior and Portfolio Selection Problem in Malaysia using Decision Making Model

Lam Weng Siew, Saiful Hafizah Hj. Jaaman and Hamizun bin Ismail

6-10

Fatty acids and Phenols Involved in Resistance of Oil Palm to *Ganoderma boninense*

Rozlianah, F.S., Jualang, A.G. and Chong, K.P.

11-16

An Empirical Study on the Characteristics of High Risk Aversion Behavior in Portfolio Decision Making using Regression Model

Lam Weng Siew, Saiful Hafizah Hj. Jaaman and Hamizun bin Ismail


17-20

An Integrated Decision Support System (DSS) for the Management of Sustainable Mariculture in Indonesia

Hermawan S. and Syafrani

21-27

The Best Scenario of Carbon Capturing Storage and Utilization (CCSU) in Malaysia



An Integrated Decision Support System (DSS) for the Management of Sustainable Mariculture in Indonesia

¹Hermawan S. and ²Syafrani

¹Civil Engineering, Faculty of Technic, Lancang Kuning University, Pekanbaru Riau, Indonesia.

²Faculty of Agriculture, Lancang Kuning University, Pekanbaru Riau, Indonesia.

ARTICLE INFO

Article history:

Received 12 February 2015

Accepted 1 March 2015

Available online 28 March 2015

Keywords:

decision support system, mariculture, sustainability, Indonesia

ABSTRACT

Indonesia has great potential for aquaculture development but its expansion is often insufficiently focused in environmentally sustainable practice. For this reason, regarding an ecosystem approach to aquaculture (EAA), the aim of this paper is to demonstrate the application of decision support system for the management of sustainable floating net cage mariculture (SYSMAR DSS) in three selected regions in Indonesia: Talise Island, Galang Island and Ekas Bay. Through GIS spatial planning tools, SYSMAR DSS is used to perform site selection and assesses production and ecological carrying capacity. Economic analysis provides vital information on 18 cases, focusing on the economic viability of Tiger Grouper (*Epinephelus fuscoguttatus*), Humpback Grouper (*Cromoliptes altivelis*) and Leopard Coral Grouper (*Plectropomus leopardus*) which utilizes various feed types and production scales. SYSMAR DSS shows that only Galang Island provides a substantial area with a suited area of about 12,940 ha. The estimated maximum production carrying capacity is 51 – 366 tons/farm and ecological carrying capacity is in the range 18,393 – 21,727 tons/year/suitable area. After a 5-year projection period, the economic evaluation highlighted that all studied culture developments are economically viable. These results confirm the SYSMAR DSS is able to determine potential sites which comply with environmental sustainability and socio-economic criteria.

© 2015 AENSI Publisher All rights reserved.

To Cite This Article: Hermawan S. and Syafrani., An Integrated Decision Support System (DSS) for the Management of Sustainable Mariculture in Indonesia. *Adv. Environ. Biol.*, 9(7), 21-27, 2015

INTRODUCTION

As the stocks of wild fish decline worldwide and the human population increases, aquaculture is becoming one of the quickest increasing food production sectors and also important as protein food supplier to the world population [1,2,3]. On the other hand as a result of the increase in fish production and seafood mariculture, there is a growing concern about the impacts of such activities on the environment. Therefore, it is vital to improve aquaculture technology and to develop management tools that address the need for an eco-friendly production process and the concerns regarding food safety [4].

In 2010, an ecosystem approach to aquaculture (EAA) was introduced by FAO to promote sustainable development, equity, and the resilience of interlinked socio-ecological systems. Farhan and Lim [5] recommended the use of Decision Support Systems (DSSs) to meet the flexibility of dynamic environments. In this paper the application of a DSS under development at the Research and Technology Centre Westcoast of the University of Kiel for the sustainable environmental and socio-economic management of floating net cage (SYSMAR) is assessed. The DSS utilizes high resolution hydrodynamic information concerning water depth, current velocities and wave heights obtained from hydrodynamic models. Through GIS as spatial planning tools, SYSMAR DSS is able to assist in estimating site selection, determine different types of carrying capacities (CC) such as production CC, as well as ecological CC to guarantee sustainable environmental development. Furthermore, a method for the economic analysis of mariculture investment is proposed. Financial indicators to enable an adequate economic assessment were identified [6,7,4].

The investigations were carried out at three priority sites in Indonesia namely Talise Island located in the northern most tip of Sulawesi; Galang Island which is part of the Riau Archipelago located opposite of

Corresponding Author: Hermawan S., Civil Engineering, Faculty of Technic, Lancang Kuning University, Pekanbaru Riau, Indonesia.

Singapore and Ekas Bay located south of the Lombok Island (see Figure 1). The development focuses on the most common high value finfish species nurtured in FNC in Indonesia [8,9,10,11]. Thus, Tiger grouper (*Epinephelus fuscoguttatus*), Humpback grouper (*Cromoliptes altivelis*), and Leopard Coral grouper (*Plectropomus leopardus*) are considered.




Fig. 1: Research Areas.

MATERIAL AND METHOD

Hydrodynamic and wave models:

In order to provide flow and wave information, hydrodynamic models for the three selected regions were set up using the open source Delft3D modeling suite. Curvilinear grids covered the region with increasing resolutions of up to 40 meters in the areas of interest as indicated in Figure 1. Bathymetric information was taken from the GEBCO database [12]. Additional bathymetric information for the near shore was provided by the national surveying agency [13]. Tidal driving forces were specified along the open model boundaries in the form of astronomical constituents [14]. Wave models were set up using maximum wind magnitudes and directions for the period 2005 to 2009 from the NCEP/NCAR reanalysis database [15].

Site Selection:

A comprehensive site suitability and capability map which integrates all of the selected criteria was edited using thirty two parameters identified in Table 1. In order to assess the relative impacts of each category of parameters (e.g. physical, chemical, and ICZM) on the DSS result, the suitability maps of the three applied categories are presented. The parameters considered in SYSMAR were taken from FAO, Cross and Kingzett, Kapetsky and Agullar-Manjarrez and Szuster and Albasri [16, 17, 18, 19].

Production Carrying Capacity:

Production carrying capacities have been defined by the emission of particulate carbon governed by the amount of waste and physical characteristics, and deposited beneath FNC farms [20]. A simplified footprint approach considers particle settling velocities, carbon flux, current velocities, water depth and dispersion constants to provide an approximation of the carbon deposition footprint and derived deposition rates around the farm as described in Gilibrand and Van der Wulp [21, 6]. Gilibrand [21] identified a breakdown rate of $0.7 \text{ kg C m}^{-2} \text{ y}^{-1}$, which is equivalent to $1.9 \text{ g C m}^{-2} \text{ d}^{-1}$ and adopted by Rachmansyah [22], while Krost [23] proposed a slightly more conservative threshold value of about 0.5 to $2 \text{ g C m}^{-2} \text{ d}^{-1}$ on the basis of measurements carried out in fish farms in Indonesia. Thus in order to ensure practical sustainability in Indonesia, we adopt the threshold value criteria of $1 - 2 \text{ g C m}^{-2} \text{ d}^{-1}$ for determining local/production carrying capacity.

Ecological Carrying Capacity:

In this paper, the ecological carrying capacity for FNC finfish farms within a domain is set to be equivalent to the production rates of total dissolved nitrogen (TDN) which do not contribute more than 1% of the TDN flux of the domain [24,25,6].

Economic Analysis of Mariculture Investments:

Economic decision tools in mariculture aim to assist farmers, potential investors, and decision makers (stakeholders) in understanding the economic requirements, costs and benefits, and risks involved in production.

Economic sustainability is maintained when environmentally sustainable production rates remain profitable [26]. In this paper, the discount cash flow analysis methodologies facilitate the justification of an investment in cases in which the net present value (NPV) is greater than 0, the benefit cost ratio (BCR) greater than 1 and the internal rate of return (IRR) greater than each benefit cost value [27,28]. Required information of the main economic properties of grouper at Galang Island was collected from various institutions in 2012 and 2013 as shown in Table 2 (1 € = 12.500 IDR).

Table 1: Site selection of groupers grown in floating net cages.

Description	Parameters	Indicators	Units	Unsuitable	Allowable	Optimal
Physical Process	Min. water depth	water depth	m	< 6	≥ 6	> 8
	Max. mooring	water depth	m	> 25	≤ 25	< 20
	Flushing	mean current	m/s	<0.01	≥0.01	0.2 – 0.5
	Currents	mean current	m/s	> 1	≤ 1	0.2 – 0.5
	Exposure to waves	significant wave max wind	m	> 1	≤ 1	< 0.6
	Exposure to wind	speed	m/s	> 15	≤ 15	< 10
	Water temperature	water temperature	°C	<20 or >35	20 – 35	27 – 31
	Salinity	salinity	ppm	<15 or >35	15 – 35	26 – 31
	Dissolved oxygen	dissolved oxygen	mgO ₂ /l	< 4	≥ 4	> 5
	Acid-base balance	pH	-log(H ⁺)	<6 or >8.5	6 – 7.8	7.8 – 8.5
Water Quality	Water transparency	Secchi depth	m	< 2	≥ 2	> 4
	Turbidity	suspended matter	mg/l	> 10	≤ 10	< 5
	Ammonium	ammonium	mg NH ₄ -N/l	> 1	≤ 1	< 0.5
	Nitrate	nitrate	mg NO ₃ -N/l	> 200	≤ 200	< 200
	Nitrite	nitrite	mg NO ₂ -N/l	> 4	≤ 4	< 4
	Phosphate	total phosphate	mg P/l	> 70	≤ 70	< 70
	Villages	thematic map	m	< 200	≥ 200	> 500
	Towns	thematic map	m	< 200	≥ 200	> 500
	Cities	thematic map	m	< 200	≥ 200	> 500
	Harbours	thematic map	m	< 200	≥ 200	> 500
ICZM	Industry	thematic map	m	< 200	≥ 200	> 500
	Tourism	thematic map	m	< 200	≥ 200	> 500
	Streams	thematic map	m	< 200	≥ 200	> 500
	Rivers	thematic map	m	< 200	≥ 200	> 500
	Erosive shoreline	thematic map	m	< 200	≥ 200	> 500
	Semi intensive hatcheries	thematic map	m	< 200	≥ 200	> 500
	Intensive hatcheries	thematic map	m	< 200	≥ 200	> 500
	Ponds	thematic map	m	< 200	≥ 200	> 500
	Sewage discharges	thematic map	m	< 200	≥ 200	> 500
	Traffic lanes	thematic map	m	< 200	≥ 200	> 500
Environmentally protected area	Coastal usage	thematic map	m	< 200	≥ 200	> 500
	Environmentally protected area	thematic map	m	< 200	≥ 200	> 500

Table 2: Economic properties of grouper cultures at Galang Island

Economic Properties		10 FNC	600 FNC
Investment , including home base, guard house, storage,	Floating net cages 3x3x3m	11,908€	504,000€
FNC, operational equipment, electric , contingency, etc	Lifespan (years)	5	
<i>Indonesian Bank, 2012</i>	Discount rate (%)	13	
<i>Progressive Tax , Law No.17, 2010[29]</i>	Tax rate (%)	25	
Food conversion ratio[30,6]	Trash fish	7.78	
	Pellets	2.64	
Grow out period (day) [6]	Tiger Grouper	300	
	Humpback Grouper	500	
	Leopard Coral Grouper	180	
Seed price(€/kg)[31]	Tiger Grouper	0.6	
	Humpback Grouper	0.96	
	Leopard Coral Grouper	2.72	
Feed price(€/kg)[31]	Trash fish	0.32	
	Pellets	1.04	
Production (kg)	Tiger Grouper	3,285	197,100
	Humpback Grouper	1,380	87,782
	Leopard Coral Grouper	5,475	328,500
Commodity Value (€/kg)[31,32]	Tiger Grouper	11.20€	
	Humpback Grouper	28.00€	
	Leopard Coral Grouper	21.60€	
Wages (€/year)	Total wages	4,612€	44,688€

RESULTS AND DISCUSSION

Site Selection:

After classifying, the physical information from hydrodynamic and wave numerical models give an insight into the spatial distribution of areas which are suited for the development of floating net cage mariculture. The first location for the application of SYSMAR DSS site selection is in the vicinity of Talise Island. The concluding outcome of the wave parameter analysis showed that all the area is defined as unsuitable (see Figure 3a). As can be seen in Figure 3b, the final result of site selection in Ekas Bay shows no potentially suitable area for development of floating net cages (FNC) grouper culture. It is affected by incompatible wave height

parameter, current and flushing which are considered too weak or low for FNC activities. Conditions are more feasible in Galang Island. The final result of the suitability map from all parameters in the vicinity of the Galang Island can be seen in Figure 3c. It shows about 12,940 Ha (40.3%) of the seawater area in Galang Island are interpreted as suited area. The areas are spread over the vicinity of the Galang Island. Regarding the results of site selection carried out on three areas, we conclude that Galang Island is a suitable location for further development of FNC grouper culture in Indonesia.

Production Carrying Capacity:

There is a large area available for the development of FNC grouper culture in Galang Island. The SYSMAR DSS is applied to recognize the best locations for a limited number of farms. The selection is carried out for all potential farms based on the production carrying capacity. Thus, suitable and potential locations for all farms with a minimum distance of 500 meters between individual farms were selected for each of the locations as shown in Figure 3d. We consider large scale farms (over 100 cages) made up of cages with the size of 3m x 3m x 3m, with a distance between cages of 1 m [33,34].




Fig. 2: Results of site selection for Talise Island (a), Ekas Bay (b), Galang Island (c) and Production carrying capacity including proposed potential farm locations as red dots (d).

In Galang Island, prediction based on maximum deposition and feeding with trash fish indicated that about 51 – 125 tons/year/farm can be produced, but when feeding with mixed trash fish and pellet or feeding just pellet about 73 to 196 tons/year/farm and 115 to 366 tons/year/farm, respectively can be sustained (see Table 3). Discrepancies of the production carrying capacity are obtained by feeding type and also physical models which are influenced by currents and maximum water depth parameters as dominant factors and responsible for the indicated ranges.

Ecological Carrying Capacity:

The ecological carrying capacity for FNC finfish farms within a domain is proposed to be equivalent to emission rates of total dissolved nitrogen (TDN) not exceeding 1% of the TDN flux of the suitable domain. The maximum daily TDN load is calculated with respect to the flushing rate and TDN background concentration of the suited region, which is recorded in the order of 0.31 mg N l^{-1} [35]. As can be seen in Table 3, the ecological carrying capacities are in the range of 18,393 - 21,727 tons per year in the vicinity of Galang Island.

Since the estimation of the total maximum production CC does not exceed the ecological carrying capacity [36], the estimation of the maximum allowable production in the vicinity of Galang Island is 21,727 t/a. Regarding the estimation of production carrying capacity in the vicinity of Galang Island, this total production is achieved by 206 fish farms with an estimated production CC in the range of 32.5 t/a/f to 366 t/a/f.

Economic Analysis of the SYSMAR DSS:

The aim of the Indonesian government is to expand fish farming activities, especially through small family owned businesses. Therefore, it is necessary to consider small scale farms (10 cages or less) and large scale farms (100 cages or more). Table 4 presents a ranking of the 18 cases studied of 10 cages and 600 cages

according to the indicators of financial viability of FNC grouper culture projects in Galang Island. The project ranking shows that the highest net present value (NPV) amounts to 9,961 million Euros that the highest internal rate of return (IRR) amounts to 590% and that the payback period is shorter than 1 year in the case of a large farm of 600 cages in which the variety Leopard Coral Grouper is fed trash fish. On the other hand, the lowest NPV is obtained from a small farm of 10 cages in which the variety Tiger Grouper is fed pellets (NPV: 34,089 €, IRR 107% and payback less than 1 year).

Table 3: Results of the estimation of site selection and carrying capacity, A) Site Selection, B) Production CC based on the dissipation of particulate organic matter in the vicinity of a particular farm, C) Ecological CC based on the maximum POM load and TDN surplus.

Type of Carrying Capacity	Feedstock	Grouper species		
		TG	HG	LG
Site Selection Ha			12,940	
Production Carrying Capacity: Deposition threshold 1g C m ⁻² d ⁻¹				
t/a/fish farm	TF ^a	0.5-59	0.5-85	0.5-51
	P ^b	0.5-158	2-400	0.5-115.5
	M ^c	0.5-90	0.5-153	0.5-73.5
Deposition threshold 2g C m ⁻² d ⁻¹				
t/a/fish farm	TF	0.5-155.5	1-277	0.5-125.5
	P	3-624	20-1000	2-366
	M	1-268	2.5-650	1-196
Ecological Carrying Capacity ^d based on a limitation by particulate organic matter:				
t/a/entire domain	TF	3,959-9,972	5,371-15,503	3,493-8,410
	P	10,478-35,993	22,162-128,071	7,942-23,795
	M	6,014-16,729	9,152-33,467	5,308-13,050
based on a limitation by Total Dissolved Nitrogen:				
t/a/entire domain	TF	18,393	18,393	18,393
	P	21,727	21,727	21,727
	M	20,116	20,116	20,166

Remarks: ^a trash fish, ^b Pellet, ^c Mix 70% trash fish and 30% pellet ^d The calculation considers an acceptable bed load of 1-2g organic carbon m⁻²d⁻¹

Surprisingly, the results for the 18 cases show strongly positive levels of NPV, very large values of internal rate of return, well above the discount rate value of 13% and an extremely short payback period below one year. It is apparent that the economic analysis for the sites in Galang Island clearly indicates the economic feasibility of FNC grouper culture projects.

The research shows that FNC finfish culture developments are economically viable as a whole, because after a 5-year projection period, positive cumulative cash flow and net present value, internal rate of return at rates above the bank rates, and a payback period far below the 5 year projected lifetime of the project are evident.

The use and implementation of a DSS for sustainable aquaculture development in Indonesia, with respect to the EAA concept of various carrying capacities which was introduced by FAO in 2010, is very complex and a number of concerns should be taken into account. We find that the dissimilarity of the definitions of carrying capacity in the different contexts, along with its development is a complex problem. On the other hand, this DSS is able to present the integration of all key components of detailed site selection, determination of carrying capacity along with an economic appraisal.

Table 4: Ranking of economic analysis of the FNC finfish culture.

No.	Type of Floating Net Cages Grouper	NPV (1.000 €)	IRR (%)	PP year
	600 Cages			
1	Leopard Coral Grouper feed with trash fishes	9,961	590	0.19
2	Leopard Coral Grouper feed with trash fishes and pellets	9,837	583	0.17
3	Leopard Coral Grouper feed with pellets	9,739	577	0.17
4	Humpback Grouper feed with pellets	4,062	257	0.39
5	Humpback Grouper feed with trash fishes	4,025	255	0.39
6	Humpback Grouper feed with trash fishes and pellets	3,994	253	0.39
7	Tiger Grouper feed with trash fishes	3,159	204	0.49
8	Tiger Grouper feed with trash fishes and pellets	3,054	200	0.50
9	Tiger Grouper 600 cages feed with pellets	2,996	197	0.51
	10 cages			
10	Leopard Coral Grouper feed with trash fishes	150	387	0.26
11	Leopard Coral Grouper feed with trash fishes and pellets	148	382	0.26
12	Leopard Coral Grouper feed with pellets	146	378	0.26
13	Humpback Grouper feed with trash fishes	51	149	0.66
14	Humpback Grouper feed with trash fishes and pellets	51	148	0.67
15	Humpback Grouper feed with pellets	50	147	0.67
16	Tiger Grouper feed with trash fishes	36	112	0.87
17	Tiger Grouper feed with trash fishes and pellets	35	109	0.89
18	Tiger Grouper feed with pellets	34	107	0.91

ACKNOWLEDGEMENTS

I would like to show my gratitude to Rector of Lancang Kuning University Pekanbaru Indonesia.

REFERENCES

- [1] Rückert S., S. Klimpel, S. Al-Quraishy, H. Mehlhorn, H.W. Palm, 2009. Transmission of Fish Parasites Into Grouper Mariculture (Serranidae: Epinephelus coioides (Hamilton, 1822)) in Lampung Bay, Indonesia. *Parasitology Research*, 104(3): 523-532.
- [2] FAO., 2007. Table of World aquaculture of fish, crustaceans, mollusks, etc., by principal producers.ftp://ftp.fao.org/fi/stat/summary/summ_07/default.htm. Accessed 15 August 2010.
- [3] Staniford, D., 2002. Sea cage fish farming: an evaluation of environmental and public health aspects (the five fundamental flaw of sea cage fish farming). Paper presented by Don Staniford at the European Parliaments Committee on Fisheries public hearing on "Aquaculture in the European Union: Present Situation and Future Prospects", 1st October 2002. http://www.europarl.eu.int/hearings/20021001/pech/programme_en.pdfhttp://www.europarl.eu.int/committees/pech_home.htm. Accessed 24 May 2011.
- [4] Hermawan, S., S.A. van der Wulp, K.R. Niederndorfer, K.H. Runte, R. Mayerle, 2012. System for the Sustainable Management of Floating Net Cage Mariculture (SYSMAR). An Application to Three Selected Regions in Indonesia. Proceeding of Marcoastecos Conference in Albania.
- [5] Farhan, A.R., S. Lim, 2010. Integrated Coastal Zone Management Towards Indonesia Global Ocean Observing System (INA-GOOS): review and recommendation, *Ocean & Coastal Management*, 52: 421-427.
- [6] Wulp, S.A., Van der, K.R. Niederndorfer, K.J. Hesse, K.H. Runte, R. Mayerle, A. Hanafi, 2010. Sustainable Environmental Management for Tropical Floating Net Cage Mariculture, A Modeling Approach, XVIIth World Congress of the International Commission of Agricultural Engineering (CIGR). Quebec City, Canada.
- [7] Mayerle, R., A. Hanafi, K.J. Hesse, S.A. Wulp, Van der, K.R. Niederndorfer, K.H. Runte, N. Ladwig, A. Giri, F. Kleinfeld, K. Sugama, 2011. Verbundprojekt: WTZ Indonesien: SYSMAR - Integriertes System für das Management einer ökologisch und sozio-ökonomisch nachhaltigen Marikultur in Indonesien; Teilproject 1. Retrieved from BMBF-Verbundvorhaben data base. (FKZ 03F0469A)
- [8] Heemstra, P.C., J.E. Randal, 1993. FAO Species Catalogue. An annotated and illustrated catalogue of the grouper, rockcod, hid, coral grouper and lyretail species, Food and Agriculture of the United Nations.<ftp://ftp.fao.org/docrep/fao/009/t0540e/t0540e00.pdf>. Accessed 7 April 2010.
- [9] Sim, S.Y., M.A. Rimmer, J.A. Toledo, K. Sugama, I. Rumengan, K.C. Williams, M.J. Philips, 2005. A Practical Guide to Feeds and Feed Management for Cultured Groupers. NACA, Bangkok. Thailand, 18.
- [10] Sadovy, Y.J., T.J. Donaldson, T.R. Graham, F. McGilvray, G.J. Muldoon, M.J. Phillips, M.A. Rimmer, A. Smith, B. Yeeting, 2003. *While Stocks Last: The Live Reef Food Fish Trade*. Pacific Studies Series. Asian Development Bank: Manila, 147.
- [11] Sadovy, Y., C. Thierry, J.H. Choat, A.S. Cabanban, 2008. *Cromileptes altivelis*. In: IUCN 2012. IUCN Red List of Threatened Species. Version 2012.2. <www.iucnredlist.org>. <http://www.iucnredlist.org/details/39774/0>. Accessed 15 December 2012.
- [12] IOC/IHO/BODC, 2003. General Bathymetric Chart of the Oceans. British Oceanographic Data Centre, Liverpool, U.K. http://www.gebco.net/data_and_products/gebco_digital_atlas/. Accessed 4 May 2010.
- [13] Bakosurtanal (Badan Koordinasi Survey dan Pemetaan Nasional), 2011. National Coordinator for Survey and Mapping Agency, Indonesia. Nautical chart number, 42: 262-334.
- [14] Egbert, G.D., S.Y. Erofeeva, 2002. Efficient Inverse Modeling of Barotropic Ocean Tides, *J. Atmos. Tide* extracted from Total Model Driver. *Oceanic Technol*, 19(2): 183-204. http://www.esr.org/polar_tide_models/Model_TPXO62.html#EgbertErofeeva_2002. Accessed 10 July 2010.
- [15] NOAA/OAR/ESRLPSD, 2009. NCEP/NCAR Reanalysis 2 data. <http://www.cdc.noaa.gov/> Global six hourly reanalysis data with the resolution 1.87 degrees (192 x 94 grid) for wind and sea level pressure. Accessed 4 May 2010.
- [16] FAO., 1989. Site Selection Criteria for Marine Finfish Net Cage Culture in Asia, UNDP/FAO Regional Sea farming Development and Demonstration Project, Network of Aquaculture Centers in Asia. FAO Documentation NACA-SF/WP/89/13.
- [17] Cross, S.F., B.C. Kingzett, 1992. Biophysical Criteria for Shellfish Culture in British Columbia : A Site Capability Evaluation System. Aquametrix Research. Ltd. Sidney. B.C., 61.

- [18] Kapetsky, J.M., J. Aguilar-Manjarrez, 2007. Geographic Information Systems, Remote Sensing and Mapping for The Development and Management of Marine Aquaculture. Rome: FAO Fisheries and Aquaculture Department, 141.
- [19] Szuster, W.B., H. Albasri, 2010. Site Selection for Grouper Mariculture in Indonesia, *International Journal of Fisheries and Aquaculture*, 2(3): 87-92.
- [20] Gowen, R.J., N.B. Bradbury, J.R. Brown, 1989. The Use of Simple Models in Assessing Two of the Interactions Between Fish Farming and the Marine Environment, in: De Pauw, N. *et al.* (Ed.). *Aquaculture: a biotechnology in progress*, 1: 1071-1080.
- [21] Gilibrand, P.A., M.J. Gubbins, C. Greathead, I.M. Davies, 2002. Scottish Executive Locational Guidelines for Fish Farming: Predicted Levels of Nutrient Enhancement and Benthic Impact. Scottish fisheries research report (63) 2002. Aberdeen: Fisheries research service, 52.
- [22] Rachmansyah., 2004. Carrying Capacity Analysis of Awerange Bay of Milkfish Culture Development in Floating Net Cages. PhD dissertation. In Indonesia Language. Bogor Agricultural University, 274.
- [23] Krost, P., unpubl.data, 2007. Sediments carrying capacity of organic pollution and cumulative effects by fish farming in a tidally influenced region in Riau region, Indonesia. SPICE Cluster 3.2 Verbundprojekt Indonesien: Entwicklung einer Systemlösung für ein nachhaltiges Management lebender Ressourcen (Aquakultur) BMBF-Verbundvorhaben FKZ 03F0393A.
- [24] Weston, D.P., 1986. The Environmental Effects of Floating Mariculture in Puget Sound. Prepared by the University of Washington, School of Oceanography for the Washington Department of Fisheries and Ecology, 148.
- [25] GESAMP, 2001. (IMO/FAO/UNESCO-IOC/WMO/WHO/IAEA/UN/UNEP Joint Group of Experts on the Scientific Aspects of Marine Environmental Protection). Planning and management for sustainable coastal aquaculture development. Report and Studies No. 68. Food and Agriculture Organization of The United Nations. <ftp://ftp.fao.org/docrep/fao/007/y1818e/y1818e00.pdf>. Accessed 16 January 2011.
- [26] Johnston, B., T. Pickering, 2003. The Economics of Aquaculture in Comparison With Other Rural Development Opportunities In Pacific Island Countries, Marine Studies Programme The University of the South Pacific, Queensland, 24.
- [27] Jacob, H., 1969. *Allgemeine Betriebswirtschaftslehre*. Verlag Gabler Wiesbaden Betriebswirtschaftlicher, 964.
- [28] Afero, F., S. Miao, A.A. Perez, 2010. Economic Analysis of Tiger Grouper *Epinephelus Fuscoguttatus* And Humpback Grouper *Cromoliptes Altivelis* Commercial Cage Culture In Indonesia, *Aquaculture Int* 18: 725 – 739. [26] Act no. 17., (2010). Undang – Undang Republik Indonesia Nomor 17 Tahun 2010 tentang Tarif efektif Pph Badan (Progressive tax for Indonesia domestic agency). <http://www.bppk.depkeu.go.id/webpajak/index.php/artikel/opini-kita-pph/1217-tarif-efektif-pph-badan> Accessed 10 January 2011.
- [29] Act no. 17., 2010. Progressive Tax for Indonesia Domestic Agency. In Indonesian language. <http://www.bppk.depkeu.go.id/webpajak/index.php/artikel/opini-kita-pph/1217-tarif-efektif-pph-badan>. Accessed 25 March 2012.
- [30] Alongi, D.M., A.D. McKinnon, R. Brinkman, L.A. Trott, M.C. Undu, M. Rachmansyah, 2009. The Fate of Organic Matter Derived From Small-Scale Fish Cage Aquaculture In Coastal Waters of Sulawesi And Sumatra Indonesia, *Aquaculture* 295(1-2): 60-75.
- [31] Ismi, S., 2012. Personal communication. GRIM (Gondol research institute for Mariculture Indonesia), Center for aquaculture research and development, Ministry of Marine Affairs and Fisheries. <http://www.rca-prpb.com/content.php>
- [32] Ministry of marine affairs and fisheries (MMAF) Indonesia, 2013. Information price of Grouper in June 2013. In Indonesian Language. <http://wartaekonomi.co.id/berita11314/budidaya-laut-peluang-usaha-berprospek-cerah.html>
- [33] Kongkeo, H., C. Wayne, M. Murdjani, P. Bunliptanon, T. Chien, 2010. Current Practices of Marine Finfish Cage Culture in China, Indonesia, Thailand and Vietnam. *Aquaculture Asia Magazine*, XV(2): 32-40.
- [34] Wong, M., M.A. Barbeau, R.A. Aiken, 1999. Intertidal Invertebrate Population Density and Diversity: Does Salmon Aquaculture Play a Role. *Environment Canada, Occ. Rep.12* : (89-100). Proceeding 3rd Bay of Fundy Workshop.
- [35] SPICE (Science for the Protection of the Indonesian Coastal Environments), unpubl.data, 2006. Development of a Decision Support System for Sustainable Environment Management of Mariculture in Indonesia. Techn. Report.
- [36] Byron, C.J., B.A. Costa-Pierce, 2010. Carrying Capacity Tolls For Use in the Implementation for an Ecosystem Approach to Aquaculture, Presented at the FAO Expert Workshop on Aquaculture Site Selection And Carrying Capacity Estimates For Inland And Coastal Water Bodies, Institute of Aquaculture, University Stirling, Stirling, U.K.